

DTIC FILE COPY

2

AD-A218 471

ABILITY DISTRIBUTIONS

PATTERN PROBABILITIES AND QUASIDENSITIES

Michael V. Levine

Model Based Measurement Laboratory
University of Illinois
210 Education Building
Champaign, IL 61820

December 1989

DTIC
ELECTE
FEB 26 1990
S E D
CO E D

Prepared under Contract No. N00014-83K-0397, NR 150-518
and No. N00014-86K-0482, NR 4421546.

Sponsored by the Cognitive Science Program
Office of Naval Research.

Approved for public release: distribution unlimited.
Reproduction in whole or in part is permitted for
any purpose of the United States Government.

75 32 26 002

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release: distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Measurement Series 89-2			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Michael V. Levine Model Based Measurement Lab.		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Cognitive Science Program Office of Naval Research		
6c. ADDRESS (City, State, and ZIP Code) University of Illinois 210 Education Building, 1310 S. Sixth St. Champaign, IL 61820			7b. ADDRESS (City, State, and ZIP Code) Code 1142PT 860 North Quincy St. Arlington, VA 22217-5000		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-83-K-0397 N00014-86-K-0482		
8c. ADDRESS (City, State, and ZIP Code)			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 61153N	PROJECT NO. RR04204	TASK NO. RR04204-01
					WORK UNIT ACCESSION NO. NR 150-518 NR 4421546
11. TITLE (Include Security Classification) Ability Distributions, Pattern Probabilities, and Quasidensities					
12. PERSONAL AUTHOR(S) Levine, Michael V.					
13a. TYPE OF REPORT Final Report		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) 1989, December	
15. PAGE COUNT 40					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Item response theory, latent trait theory, latent class theory, multiple choice tests, density estimation, distribution estimation, ability distributions, (continued)		
05	09				
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The quasidensity is a useful surrogate for the ability density in item response theory analyses of multiple choice tests. Like the ability density, it can be used to calculate the probability of sampling an examinee with a specified pattern of responses. Sometimes it is preferable to the density because it is continuous (densities need not be continuous), it is unique (two very different densities can give exactly the same pattern probabilities and all other expected values of random variables that are functions of item responses), it always exists (a discontinuous ability distributions has a quasidensity, but it does not have a density). Some large sample results are proven for quasidensity estimation. It is shown that the maximum likelihood quasidensity estimate (mle) is strongly consistent. The asymptotic distribution of the mle is derived. Some new results on the relation between latent class and latent trait models are also presented. It is shown that every item response model with a smooth density and smooth item response-(continued)					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Charles Davis			22b. TELEPHONE (Include Area Code) 202-696-4046		22c. OFFICE SYMBOL ONR 1142CS

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

18. continued ...


nonparametric density estimation, foundations of measurement, formula score theory, Rasch model, logistic models, consistent estimation, asymptotic distribution, quasidensities, large sample theory, consistency, asymptotic normality.

19. continued ...

functions is isomorphic to the latent class model obtained with the same item response functions and a discrete ability distribution. An upper bound for the number of points is derived.

(122)

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



UNCLASSIFIED

ABILITY DISTRIBUTIONS
PATTERN PROBABILITIES AND QUASIDENSITIES

Michael V. Levine

Model Based Measurement Laboratory
University of Illinois
210 Education Building
Champaign, IL 61820

December 1989

Prepared under Contract No. N00014-83K-0397, NR 150-518
and No. N00014-86K-0482, NR 4421546.

Sponsored by the Cognitive Science Program
Office of Naval Research.

Approved for public release: distribution unlimited.
Reproduction in whole or in part is permitted for
any purpose of the United States Government.

Preface

This paper was originally intended as a report on some large sample results for quasidensity and pattern probability estimation, results that provide a foundation for our methods for ability distribution estimation and item response function estimation. While attempting to determine the generality of one of the results, I happened upon a promising new line of research. The new methods and results (Section Three) admittedly haven't been integrated into this report very well. The main result, relating models with a continuum of abilities to finite models, seems at least as important as the large sample results. (It asserts that every item response model with a smooth ability distribution and smooth item response functions is isomorphic to a latent class model obtained by replacing the ability distribution with a discrete distribution.) The reader primarily interested in the generality of latent class models may wish to skim Sections One and Two for notation and then read Section Three. Two separate papers eventually will be prepared for publication.

ABILITY DISTRIBUTIONS, PATTERN PROBABILITIES, AND QUASIDENSITIES

Introduction

This paper solves a problem closely related to ability distribution estimation, which is arguably the central problem of item response theory. The problem, quasidensity estimation, comes up when one needs to know the probability of sampling an examinee with a specified item response pattern from a very large pool of examinees. The situation in which item response functions are specified but nothing is known about the distribution of ability is considered in this paper.

If the ability distribution has a density, then the density can be used to calculate the probability of sampling an examinee with a specified pattern. A pattern's probability is simply the integral of the product of the pattern's likelihood function times the ability density.

It is shown that even if the ability distribution is a step function or some other distribution that doesn't have a density there is an essentially unique, continuous function that can be used in place of a density to compute pattern probabilities. The integral of the product of this function (the quasidensity) and any pattern's likelihood function is exactly equal to the pattern's probability.

When the ability distribution is unknown, estimates must be used. Quasidensity estimation is easier than ability distribution estimation because the quasidensity is identifiable, but the ability distribution is not. Many different ability distributions will fit large samples equally well (Levine, 1989). By contrast, the maximum likelihood quasidensity estimate is unique (Section II.2, below).

Quasidensity estimation is related to ability distribution estimation in two ways. First, under general conditions the indefinite integral of the

quasidensity equals or closely approximates the cumulative distribution of ability. Second, the methods of quasidensity estimation have been generalized to obtain a nonparametric theory for ability density estimation.

As others (Lord, 1970; Samejima, 1981) have observed, item response function estimation is closely related to ability distribution estimation. The results in this paper are central to our current work on nonparametric item response function and option response function estimation.

A quasidensity estimation theory is developed in this paper. The quasidensity is represented as a linear combination of orthogonal functions. The set of linear combination coefficients is shown to be convex and compact. It is shown that the maximum likelihood estimate of the coefficients is strongly consistent. The asymptotic distribution of the coefficients is derived.

Some general results on ability distributions are also proven. For example, it is shown that for the most commonly used item response models, every ability distribution is equivalent to a distribution with only finitely many points of increase. An upper bound for the number of points of increase is obtained.

Section One

Quasidensities, Parameterizations, and Approximations

In many applications it is necessary to compute the probability of sampling an examinee with a specified item response function. For example, pattern probabilities are needed in situations in which it is important to decide whether because of cheating, language problems or an ill-advised test-taking strategy, an individual's test-taking behavior is so unlike other examinees' that his/her test score is virtually uninterpretable. With pattern probabilities a uniformly most powerful statistical test for faulty answer sheets can be computed (Levine and Drasgow, 1988).

In this section a general strategy for computing pattern probabilities is derived and discussed. We begin with some notation and a brief review of basic item response theory.

Section I.1: Terminology, Notation and Assumptions

Let $u^* = \langle u_1^*, u_2^*, \dots, u_n^* \rangle$ denote a vector of ones and zeros indicating right and wrong answers to n test items. Sampled vectors are *locally independent* relative to a random variable θ if the conditional probability of sampling pattern u^* given any value of θ can be factored and written

$$\text{Prob}(u = u^* | \theta = t) = \prod_{i=1}^n \text{Prob}(u_i = u_i^* | \theta = t)$$

where u denotes the sampled vector, u_i is its i th component and t is one of the possible values of θ .

Usually in item response theory the *item response functions* $P_i(t) = \text{Prob}(u_i = 1 | \theta = t)$ are assumed to have some specific (generally logistic) functional form with values strictly between zero and one. The results in this paper (except for the last part of Section III) assume only that the

item response functions are continuous functions with values that are strictly between zero and one.

Usually item response functions are defined over an unbounded range and applied over a finite range, typically the interval $[-3,3]$. The results in this paper only use item response function values over a finite range of abilities. Thus, throughout this paper the P_i are continuous functions with values strictly between zero and one that are defined on some finite closed interval $[c,d]$. This results in no loss of generality because the interval c can be very large and because an unbounded ability continuum can be transformed into an interval. In applications, we make the interval $[c,d]$ big enough so that the assumption that all abilities are in $[c,d]$ is plausible.

This paper is concerned with *distributions on $[c,d]$* , i.e. distributions of random variables that are between c and d with probability one. The condition that a distribution function G is a distribution on $[c,d]$ can be expressed without explicitly referring to random variables as follows: for $t < c$, $G(t) = 1 - G(d)$.

To summarize, the results to follow assume

1. local independence relative to a unidimensional random variable θ ,
2. continuous item response functions defined on an interval $[c,d]$ and taking values strictly between zero and one, and
3. $\text{Prob}(c \leq \theta \leq d) = 1$.

Section I.2: The Canonical Space and its Quasidensities

The assumption that the P_i are strictly between zero and one implies that for every pattern u^* , the *pattern likelihood function* $\ell(u^*, \cdot)$ given by

$$\ell(u^*, t) = \text{Prob}(u=u^* | \theta=t) = \prod_1^n P_i(t)^{u_i^*} [1 - P_i(t)]^{1-u_i^*}$$

will also be a continuous, positive function defined on $[c, d]$. By forming linear combinations of likelihood functions we obtain a finite dimensional real vector space which is called the *test's canonical space* (CS). Thus f is in the CS if and only if for some real constants a_ν ,

$$f(\cdot) = \sum_{\nu=1}^{2^n} a_\nu \ell(u_\nu^*, \cdot)$$

where $u_1^*, u_2^*, \dots, u_\nu^*, \dots$ is any enumeration of the 2^n possible item response patterns. Since the functions in the CS are continuous, an inner product for the CS is defined by $\langle f, g \rangle = \int_c^d f(t)g(t) dt$.

Note that since the likelihood functions may be linearly dependent, it may be possible to write a function as a linear combination of likelihood functions in several ways. The uniqueness referred to in the following result applies to functions, not vectors of coefficients (a_ν) .

I.1 There is a unique function g in the CS such that for all response patterns u^*

$$\text{Prob}(u=u^*) = \int_c^d \ell(u^*, t)g(t) dt.$$

Equivalently, for any distribution G on $[c, d]$ there is a unique g in the CS such that

$$\int_c^d \ell(u^*, t) dG(t) = \int_c^d \ell(u^*, t)g(t) dt$$

Proof: The formula

$$h = \sum_{\nu} a_{\nu} \ell_{\nu} \rightarrow E[h(\theta)] = \sum_{\nu} a_{\nu} \text{Prob}(u=u_{\nu}^*)$$

defines a linear mapping on the CS since expectation is linear. Since every linear functional defined on an inner product space that is isomorphic to a Euclidean space can be written as an inner product, there is some g in the CS such that the mapping can be written

$$\sum_{\nu} a_{\nu} \ell_{\nu}(\cdot) \rightarrow \langle \sum_{\nu} a_{\nu} \ell_{\nu}(\cdot), g \rangle .$$

In particular, if all the a 's are zero except one, then for every u_{ν}^*

$$\text{Prob}(u=u_{\nu}^*) = \langle \ell(u_{\nu}^*, \cdot), g \rangle .$$

If \bar{g} also satisfies these conditions then for all ν

$$\begin{aligned} 0 &= \langle \ell_{\nu}, g \rangle - \langle \ell_{\nu}, \bar{g} \rangle \\ &= \langle \ell_{\nu}, g - \bar{g} \rangle . \end{aligned}$$

Thus $g - \bar{g} = 0$ because no nonzero element of a vector space can be orthogonal to all of the vector space's generators. //

The function g is called the *quasidensity* of G because it functions like a density in the calculation of pattern probabilities. In fact, it can be used in place of a density to calculate the expected value of any statistic that is a function of item responses (Levine, 1989, Section 2). Although the quasidensity integrates to one, it is generally not non-negative. A discussion of their properties can be found in (Levine, 1989).

When an orthonormal basis for the CS is available the quasidensity has a simple formula which is often used.

I.2: If θ has distribution G where G is a distribution on $[c,d]$ and

$$h_0, h_1, \dots, h_J$$

is an orthonormal basis for the canonical space, then the quasidensity for G is

$$g(\cdot) = \sum_{j=0}^J E[h_j(\theta)] h_j(\cdot) .$$

Proof: Since $\{h_j\}_{j=0}^J$ is an orthonormal basis, each pattern likelihood function satisfies $\ell(u_\nu^*, t) = \sum_j \langle \ell_\nu, h_j \rangle h_j(t)$. Consequently

$$\begin{aligned} P(u = u_\nu^*) &= \int_c^d \sum_j \langle \ell_\nu, h_j \rangle h_j(t) dG(t) \\ &= \sum_j \langle \ell_\nu, h_j \rangle E[h_j(\theta)] \\ &= \sum_j \int_c^d \ell_\nu(t) h_j(t) dt E[h_j(\theta)] \\ &= \int_c^d \ell(u_\nu^*, t) \sum_j E[h_j(\theta)] h_j(t) dt . \end{aligned}$$

From uniqueness proven in I.1 it follows that $g(\cdot) = \sum_j E[h_j(\theta)] h_j(\cdot)$. //

The value of the quasidensity in studying pattern probabilities derives from the following obvious but very useful fact:

I.3: If $\{h_j\}_{j=0}^J$ is a basis for the CS and the quasidensity distribution of θ satisfies

$$g(\cdot) = \sum_0^J \pi_j h_j(\cdot)$$

then

$$\text{Prob}(u = u_\nu^*) = \sum_0^J \pi_j \langle h_j, \ell_\nu \rangle .$$

Thus, each pattern probability can also be represented as an inner product of a known vector depending only on the pattern and a vector that must be estimated.

Section I.3: Approximations and Bases

For applications, the size of J is important. For the Rasch model $(P_i(t) = [1 + e^{-(t-b_i)}]^{-1})$ J cannot be larger than the number of items, n (Levine, 1989, Section 2). On the other hand for the three parameter logistic model $(P_i(t) = c_i + (1-c_i)[1 + e^{-a_i(t-b_i)}]^{-1})$, J may be as large as 2^n . Fortunately by careful choice of the functions h_j , approximations can be obtained that have very few terms but are still accurate in one sense or another. Two examples follow.

For a first example suppose it is desirable to keep the total squared error

$$\sum_{\nu} [\text{Prob}(u = u_{\nu}^*) - \text{approximated Prob}(u = u_{\nu}^*)]^2$$

small. A basis can be obtained by analyzing the function of two variables

$$H(s, t) = \prod_{i=1}^n (P_i(s)P_i(t) + [1-P_i(s)][1-P_i(t)])$$

defined for $c \leq s, t \leq d$. By solving the functional equation

$$\int_c^d H(s, t)h(s)ds = \lambda h(t)$$

one obtains a maximal set of orthonormal functions h_j in the CS and positive constants $\lambda_0 \geq \lambda_1 \geq \dots \lambda_J > 0$ such that

$$H(s, t) = \sum_{j=0}^J \lambda_j h_j(s)h_j(t).$$

These functions can be shown to form an orthonormal basis for the CS (Levine 1989). From this fact and the identity

$$H(s, t) = \sum_{\nu=1}^{2^n} \ell(u_{\nu}^*, s)\ell(u_{\nu}^*, t)$$

it follows that the total squared error with a $K < J$ term approximation satisfies

$$\sum_{\nu} [\text{Prob}(u = u_{\nu}^*) - \sum_{j < K} \pi_j \langle \ell_{\nu}, h_j \rangle]^2 = \sum_{j \geq K} \lambda_j \langle g, h_j \rangle^2$$

$$< \lambda_K \int_c^d g^2(t) dt$$

when $\pi_j = \langle g, h_j \rangle$.

This relation is important because for all tests we have analyzed, the λ_j very rapidly decrease to zero. Typically, $K=15$ provides very accurate least squares pattern probability approximation.

To introduce a second and final example concerning the choice of a CS basis, suppose it is important to control the maximum absolute error. In addition suppose an approximation of the distribution function G is available. Then it is often possible to select an orthonormal basis h_j such that $E[h_j(\theta)]$ will be small for large j . Of course, small $E[h_j(\theta)]$ guarantees that the finite sum $\sum_{j \geq K} |E[h_j(\theta)]|$ is also small. This is important because for $\pi_j = \langle g, h_j \rangle = E[h_j(\theta)]$ the K term approximation satisfies

$$\begin{aligned} |\text{Prob}(u = u_{\nu}^*) - \sum_{j < K} \pi_j \langle \ell_{\nu}, h_j \rangle| &= \sum_{j \geq K} |\pi_j \langle \ell_{\nu}, h_j \rangle| \\ &\leq \left(\sum_{j \geq K} |\pi_j| \right)^{1/2} \left(\sum_{j \geq K} |\pi_j| \langle \ell_{\nu}, h_j \rangle^2 \right)^{1/2} \\ &\leq \left(\sum_{j \geq K} |\pi_j| \right)^{1/2} \text{Max}_{j \geq K} |\pi_j|^{1/2} \left(\sum_{j \geq K} \langle \ell_{\nu}, h_j \rangle^2 \right)^{1/2} \\ &\leq \left(\sum_{j \geq K} |\pi_j| \right)^{1/2} \text{Max}_{j \geq K} |\pi_j|^{1/2} \left(\int_c^d \ell_{\nu}^2(t) dt \right)^{1/2} \\ &\leq \text{Max}_{j \geq K} |\pi_j|^{1/2} \left(\sum_{j \geq K} |\pi_j| \right)^{1/2} (d-c)^{1/2} \text{Max}_{c \leq t \leq d} \ell_{\nu}(t) \end{aligned}$$

This paper is concerned with using maximum likelihood estimation of the π_j , from sampled patterns to obtain approximations of pattern probabilities. It will be shown that the maximum likelihood estimate for the vector π is strongly consistent and asymptotically efficient. In addition some results about the set of vectors π corresponding to distributions are proven.

Section Two

Uniqueness and Consistency of the Maximum Likelihood Estimate

In the remainder of this paper F is used to denote the unknown distribution of the ability random variable θ , $(h_j)_{j=0}^J$ is a fixed orthonormal basis for the GS, and π is the vector of expectations with j th coordinate $\pi_j = E[h_j(\theta)]$. Thus the pattern probability for u^* can be written

$$P_{\pi}(u^*) = \sum \langle \ell(u^*, \cdot), h_j \rangle \pi_j$$

or with the abbreviation $\beta_j(u^*) = \langle \ell(u^*, \cdot), h_j \rangle$, $P_{\pi}(u^*) = \beta(u^*) \cdot \pi$. If G is the distribution of θ or any other distribution on $[c, d]$ and $g_{\pi}(\cdot) = \sum \pi_j' h_j(\cdot)$ is its quasidensity then the pattern probabilities obtained by using G in place of F are given by

$$\begin{aligned} P_{\pi'}(u^*) &= \int_c^d \ell(u^*, t) dG(t) \\ &= \beta(u^*) \cdot \pi'. \end{aligned}$$

This section begins the task of estimating F 's quasidensity, or equivalently the vector π , from a sample of patterns u^* .

Section II.1: Distributions Viewed as Points in a Convex, Compact Subset of Euclidean Space

Procedures for recovering the unknown π_j from samples of observed patterns have been developed. Each requires the maximization of some continuous function defined over a set A of vectors corresponding to quasidensities

$$A = \{ \pi' \text{ in } E^{J+1} : \sum \pi_j' h_j(\cdot) \text{ is the quasidensity of at least one distribution on } [c, d] \}.$$

The set A has three properties that greatly simplify maximization:

- i. Convexity: the line segment connecting any two points in

A is also in A .

ii. Boundedness: There is a constant κ such that for all π' in A $|\pi' - \pi| = \langle \pi' - \pi, \pi' - \pi \rangle^{1/2} \leq \kappa$.

iii. Closure: Every continuous, bounded function defined on A has a maximizer in A .

The proof requires a simple result that is repeatedly used elsewhere.

II.1: The vectors $(\beta(u_\nu^*))_{\nu=1}^{2^n}$ span a $J+1$ dimensional vector space.

Equivalently, for any choice of positive constants w_ν the matrix

$$\sum_{\nu=1}^{2^N} \beta(u_\nu^*) \beta(u_\nu^*)^T w_\nu$$

is positive definite.

Proof: Let $\ell_{\nu_1}, \ell_{\nu_2} \dots \ell_{\nu_{J+1}}$ be likelihood functions forming a basis for the CS. Since $\ell_{\nu_i}(\cdot) = \sum_j \beta_j(u_{\nu_i}^*) h_j(\cdot)$, linear independence of the ℓ_{ν_i} implies linear independence of the $\beta(u_{\nu_i}^*)$. Thus the set $(\beta(u_\nu^*))_{\nu=1}^{2^n}$ of $J+1$ vectors contains $J+1$ linearly independent vectors. //

II.2: The set A of coordinates of distributions on $[c,d]$ is convex, closed and bounded.

Proof: (i) Convexity: For π^1 and π^2 in A and $0 \leq \epsilon \leq 1$ let G_1 and G_2 be distributions on $[c,d]$ with quasidensities g_{π^1} and g_{π^2} respectively. Since for any positive ϵ , $G_3 = \epsilon G_1 + (1-\epsilon)G_2$ is also a distribution on $[c,d]$ and since for any h in the CS $\int_c^d h(t) d[\epsilon G_1(t) + (1-\epsilon)G_2(t)] = \epsilon \int_c^d h(t) dG_1(t) + (1-\epsilon) \int_c^d h(t) dG_2(t)$, it follows from I.1 that $g_{\epsilon\pi^1 + (1-\epsilon)\pi^2}$ is the quasidensity of G_3 . Thus a convex combination of points in A is in A .

(ii) Boundedness: Let π' be any vector in A other than π . Let, $\bar{\pi}'$ be the intersection of the unit sphere about π and the ray from π passing through π' . Thus $|\bar{\pi}' - \pi| = 1$ and $\bar{\pi}' = [\pi' - \pi]/k + \pi$, (or equivalently, $\pi' = \pi + k(\bar{\pi}' - \pi)$) for $k = |\pi - \pi'| > 0$. Since $\pi' \cdot \beta(u^*) = P_{\pi'}(u^*)$ and $\pi \cdot \beta(u^*) = P_{\pi}(u^*)$ are probabilities,

$$0 \leq P_{\pi'}(u^*) = [\pi + k(\bar{\pi}' - \pi)] \cdot \beta(u^*) \leq 1$$

and

$$-1 \leq -P_{\pi}(u^*) \leq k(\bar{\pi}' - \pi)^T \beta(u^*) \leq 1 - P_{\pi}(u^*) \leq 1.$$

After squaring and summing over all 2^n patterns we obtain

$$0 \leq k^2 (\bar{\pi}' - \pi)^T \sum_u \beta(u^*) \beta(u^*)^T (\bar{\pi}' - \pi) \leq 2^n.$$

But in II.1, $\sum_u \beta(u^*) \beta(u^*)^T$ was shown to be positive definite. Since $|\bar{\pi}' - \pi| = 1$, the expression multiplying k^2 must be at least as large as the smallest eigenvalue of this matrix. Thus for $\kappa^2 = 2^n / [\text{smallest eigenvalue}]$, $|\bar{\pi}' - \pi| \leq \kappa$, and A is bounded.

(iii) Closure: Let $\{\pi^n\}$ be a Cauchy sequence in A . Since A is bounded, the sequence converges to some π' in E^{J+1} . To show π' is in A , let $\{G_n\}$ be any sequence of distributions on $[c, d]$ such that g_{π^n} is the quasidensity of G_n . By Helly's theorem, $\{G_n\}$ contains a subsequence $\{G_{m(n)}\}$ such that $G_{m(n)}$ converges to some distribution G on $[c, d]$ at every point of continuity of G . Since each h_j is continuous, $\int_c^d h_j(t) dG_{m(n)}(t) \rightarrow \int_c^d h_j(t) dG(t)$. Thus, by I.2 $\pi_j^{m(n)} \rightarrow \int_c^d h_j(t) dG(t)$. Since $\{\pi^{m(n)}\}$ is a subsequence of $\{\pi^n\}$, $\pi_j^{m(n)} \rightarrow \int_c^d h_j(t) dG(t) = \pi'_j$, and $g_{\pi'}$ is the quasidensity of G . //

Note that since the CS is also a metric space with distance $(g, h) =$

$\langle g-h, g-h \rangle^{1/2}$ and since $\pi' \rightarrow g_\pi$, is an isometry, it follows that the subset of the CS corresponding to quasidensities of distributions on $[c,d]$ is also convex and compact.

Section II.2: Uniqueness of the Maximum Likelihood Estimate

Consider now drawing a random sample of N patterns

$$u_1^*, u_2^*, \dots, u_a^*, \dots, u_N^*$$

and attempting to recover the quasidensity g_π by maximizing the sample likelihood function $\prod_{a=1}^N P_{\pi'}(u_a^*)$ defined for vectors π' in A or its logarithm

$$L_N(\pi') = \sum_{a=1}^N \log \pi' \cdot \beta(u_a^*) .$$

It will be shown that if the sample is large enough that L_N almost surely has a unique maximum $\hat{\pi}_N$.

II.3: With probability one $L_N(\cdot)$ eventually has a unique maximizer in A .

Proof: For each pattern u^* and vector π' in A , $P_{\pi'}(u^*) = \pi' \cdot \beta(u^*)$ is positive. Therefore, $L_N(\cdot)$ is defined and continuous on A . Since A is compact, $L_N(\cdot)$ has at least one maximizer in A . Suppose $\pi' \neq \pi''$ both maximize A . Since the line segment connecting two points of A is entirely in A , a function of one variable is defined by

$$p(\epsilon) = L_N[\pi' + \epsilon(\pi' - \pi'')]$$

for $0 \leq \epsilon \leq 1$. From the formula for L_N and the fact the π' and π'' are maximizers, p has 2 continuous derivatives, and $p(0) = p(1)$. Since the second derivative of p is the negative of a sum of squares

$$p''(\epsilon) = -\sum_{a=1}^N w_a(\epsilon) [(\pi' - \pi'') \cdot \beta(u_a^*)]^2$$

for $w_a(\epsilon) = (P_{\pi'}(u_a^*) + \epsilon [P_{\pi''}(u_a^*) - P_{\pi'}(u_a^*)])^{-2} > 0$ and $\max_{\epsilon} p(\epsilon) = p(0) = p(1)$, p must be constant. Thus, the second derivative of p evaluated at, say, $\epsilon=.5$ must be zero. However,

$$0 = -\sum_{a=1}^N w_a(.5) (\pi' - \pi'')^T \beta(u_a^*) \beta(u_a^*)^T (\pi' - \pi'')$$

implies that the N vectors $\beta(u_a^*)$ span a subspace of dimension less than $J+1$. Since each of the 2^n response patterns has positive probability of being sampled and since by II.1 the full set spans a space of dimension $J+1$, with probability one eventually a linearly independent set of $J+1$ patterns will be sampled, and the maximizer will henceforth be unique. //

Section II.3: Strong Consistency of the Maximum Likelihood Estimate

In order to study the asymptotic behavior of $\hat{\pi}_N$, the maximum likelihood estimate, it is convenient to have an open set containing A on which any L_N can be extended to a differentiable function. To this end we choose a positive number d such that if a vector x is within distance d of at least one point of A , then $x \cdot \beta(u^*) \geq 0$ for all patterns u^* .

II.4: Let $d = \inf_{u^*} \text{UNION} \{ |x - \pi'| : \pi' \text{ is in } A \text{ and } x \cdot \beta(u^*) = 0 \}$ and

$$A^+ = \{x : \text{for some } \pi' \text{ in } A, |x - \pi'| < d\}.$$

Then A^+ is an open set containing A on which the formula

$$\sum_{a=1}^N \log x \cdot \beta(u_a^*)$$

extends L_N to a differentiable function defined on A^+ .

Proof: It remains only to show d is positive. Since A is compact, for each u^* the set $\{ |x - \pi'| : x \cdot \beta(u^*) = 0 \text{ and } \pi' \text{ is in } A \}$ has a positive minimum. Thus d is the minimum of 2^n positive numbers. //

If there is some π' in A such that $P_{\pi'}(\cdot)$ assigns exactly the same probabilities to patterns as $P_{\pi}(\cdot)$ then it will not be possible to prove $\hat{\pi}_N$ converges to π . Thus the following intrinsically important result is needed.

II.5: If $\pi' \neq \pi''$ and both are in A , then for at least one pattern u^* , $P_{\pi'}(u^*) \neq P_{\pi''}(u^*)$.

Proof: If $P_{\pi'}(\cdot) = P_{\pi''}(\cdot)$, then for all u^*

$$0 = \beta(u^*) \cdot (\pi' - \pi'')$$

Since from II.1 the $\beta(u^*)$ span a $J+1$ dimensional space, the $J+1$ vector π' must equal π'' . //

Finally, strong consistency for the maximum likelihood estimate can be proven by an argument Wald used to prove the consistency in a different context (Wald, 1949).

II.6: (Strong consistency of the mle) With probability 1, $\hat{\pi}_N$ converges to π .

Proof: Using the inequality $\log x < x-1$ for $x \neq 1$ and II.5 it follows that for $\pi' \neq \pi$, $E \log P_{\pi'}(u) < E \log P_{\pi}(u)$. For if the set $D = \{ u^* : P_{\pi'}(u^*) \neq P_{\pi}(u^*) \}$ is not empty then

$$\begin{aligned} E [\log P_{\pi'}(u)] - E [\log P_{\pi}(u)] &= E \log [P_{\pi'}(u) / P_{\pi}(u)] \\ &= \sum_{u^* \text{ in } D} P_{\pi}(u^*) \log [P_{\pi'}(u^*) / P_{\pi}(u^*)] \\ &< \sum_{u^* \text{ in } D} P_{\pi}(u^*) [P_{\pi'}(u^*) / P_{\pi}(u^*) - 1] \\ &= \sum_D P_{\pi'}(u^*) - \sum_D P_{\pi}(u^*) \end{aligned}$$

$$\begin{aligned}
&= \sum_D P_{\pi'}(u^*) - [1 - \sum_{u^* \text{ not in } D} P_{\pi'}(u^*)] \\
&= 0
\end{aligned}$$

To obtain a finite open covering of some subsets of A , note that $x \cdot \beta(u^*)$ is positive for x in A^+ . For $\pi' \neq \pi$ let $s(u^*, \pi', \rho) = \sup \{\pi'' \cdot \beta(u^*) : \pi'' \text{ is in } A^+ \text{ and } |\pi' - \pi''| < \rho\}$. Since for π'' in A , π'' in A^+ and $|\pi' - \pi''| < \rho$

$$\begin{aligned}
\pi'' \cdot \beta(u^*) &= [\pi' + (\pi'' - \pi')] \cdot \beta(u^*) \\
&\leq P_{\pi'}(u^*) + |\pi'' - \pi'| |\beta(u^*)| \\
&< P_{\pi'}(u^*) + \rho |\beta(u^*)|,
\end{aligned}$$

$\log s(u^*, \pi', \rho) \leq \log P_{\pi'}(u^*) + \rho |\beta(u^*)| / P_{\pi'}(u^*)$. Consequently

$$\begin{aligned}
E \log s(u, \pi', \rho) &\leq E \log P_{\pi'}(u) + \rho E[|\beta(u)| / P_{\pi'}(u)], \\
&= E \log P_{\pi}(u) \\
&\quad - [E \log P_{\pi}(u) - E \log P_{\pi'}(u)] \\
&\quad + \rho E[|\beta(u)| / P_{\pi'}(u)]
\end{aligned}$$

and for each $\pi' \neq \pi$ in A , a positive $\rho(\pi')$ less than d can be selected so that

$$E \log s(u, \pi', \rho(\pi')) < E \log P_{\pi}(u).$$

Let B be a closed subset of A not containing π . To show that with probability one

$$\sup_{\pi'' \text{ in } B} \prod_{a=1}^N P_{\pi''}(u_a^*) / P_{\pi}(u_a^*)$$

tends to zero as N tends to infinity, consider the open covering of A formed by the sets

$$B(\pi') = \{\pi'' \text{ in } A^+ : |\pi'' - \pi'| < \rho(\pi')\}.$$

Since B is also compact $\pi'_1, \pi'_2, \dots, \pi'_m$ can be selected such that $B(\pi'_1), B(\pi'_2), \dots, B(\pi'_m)$ covers B . By the strong law of large numbers,

$$\frac{1}{N} \sum_{a=1}^N \{ \log s(u_a^*, \pi'_i, \rho(\pi'_i)) - \log P_{\pi}(u_a^*) \}$$

almost surely tends to $E \log s(u, \pi'_i, \rho(\pi'_i)) - E \log P_{\pi}(u) < 0$.

Consequently with probability one,

$$\sum_{a=1}^N \log s(u_a^*, \pi'_i, \rho(\pi'_i)) / P_{\pi}(u_a^*) \rightarrow -\infty$$

and $\lim_N \sup_{\pi'' \text{ in } B_i} \prod_{a=1}^N \pi'' \cdot \beta(u_a^*) / P_{\pi}(u_a^*) = 0$. Since

$$\begin{aligned} 0 &< \sup_{\pi'' \text{ in } B} \prod_{a=1}^N P_{\pi''}(u_a^*) \\ &\leq \sum_{i=1}^m \sup_{\pi'' \text{ in } B_i} \prod_{a=1}^N \pi'' \cdot \beta(u_a^*), \end{aligned}$$

with probability one

$$\lim_N \sup_{\pi'' \text{ in } B} \prod_{a=1}^N P_{\pi''}(u_a^*) / \prod_{a=1}^N P_{\pi}(u_a^*) = 0.$$

Finally, to show $\hat{\pi}_N$ converges almost surely to π , it is shown that for any positive ϵ , with probability one, $|\hat{\pi}_N - \pi|$ is eventually less than ϵ .

Let B be the closed set excluding π ,

$$B = \{ \pi' \text{ in } A : |\pi' - \pi| \geq \epsilon \}.$$

With probability one

$$\sup_{\pi' \text{ in } B} \prod_{a=1}^N P_{\pi'}(u_a^*) / \prod_{a=1}^N P_{\pi}(u_a^*)$$

is less than one for sufficiently large N . Thus with probability one $\hat{\pi}_N$ is eventually in the complement of B , i.e., $|\hat{\pi}_N - \pi|$ is eventually less than ϵ . //

Section III

A Dichotomy for Ability Distributions

A common starting point for studying the asymptotic distribution of maximum likelihood estimates is the family of likelihood equations

$$0 = \frac{\partial}{\partial \pi_j} L_N(\hat{\pi}_N) \quad j=0,1, \dots J .$$

It turns out that these equations are false for our current formulation of the estimation problem.

The equations are valid if $\hat{\pi}_N$ is the maximum of L_N over an open $J+1$ dimensional subset of A . But A has no $J+1$ dimensional subsets because, as shown below, it is a J dimensional subset of a $J+1$ dimensional space.

In this section A is reparameterized as a J dimensional set, i.e., the points of A are expressed as functions of J numbers. The reparameterization suggests a dichotomy of the distributions on $[c,d]$. We distinguish "regular" distributions that correspond to points in the interior of the new set of parameters for A and "irregular" distributions that correspond to boundary points. The distinction is important because in this paper the asymptotic distribution of the mle is worked out in detail only for regular distributions.

In the process of attempting to show that the irregular distributions were pathological and safe to ignore a surprising result was obtained. It was found that for the most popular item response models, every distribution is equivalent to some discrete probability distribution on at most $J+1$ points.

Section III.1: Reparameterization

Since $1 = \sum_u P_{\pi'}(u^*) = \pi' \cdot \sum_u \beta(u^*)$ every vector π' in A satisfies the equation $\pi' \cdot \bar{\beta} = 1$ for $\bar{\beta} = \sum_u \beta(u^*)$. Thus A is contained in a J dimensional subset of E^{J+1} and no point of A has an open neighborhood contained in A .

To reparameterize A let $(\bar{\beta}/|\bar{\beta}|, z^1, z^2, \dots, z^J)$ be an orthonormal basis for E^{J+1} and Z be the $(J+1) \times J$ matrix

$$Z = [z^1, z^2, \dots, z^J] .$$

Thus $x \rightarrow Z^T x$ maps the set of vectors orthogonal to $\bar{\beta}$ one-to-one onto E^J . Since $x \rightarrow Z^T x$ is also linear, it follows that for any fixed π^0 in A

$$\pi' \rightarrow Z^T(\pi' - \pi^0)$$

maps A one-to-one, onto some convex, compact subset of E^J and that each π' in A can be expressed as

$$\pi' = \pi^0 + Zs$$

for exactly one J vector s in a convex, compact subset of E^J .

Let $B = \{Z^T(\pi' - \pi^0) : \pi' \in A\}$ be the set of possible J vectors s . It will be shown that B is J dimensional so that distributions on $[c, d]$ can be classified into two non-empty sets: regular distributions with s in the interior of B and irregular distributions corresponding to vectors on the boundary of B . Before resuming the study of the distribution of the maximum likelihood estimate some facts about regular and irregular distributions will be proven.

Section III.2: Regular and Irregular Distributions

A distribution on $[c,d]$, its quasidensity $g_{\pi'}$, and its coordinate vector π' will be called *regular* if for some $\rho > 0$ the J -dimensional open ball centered at π'

$$\{\pi'' \text{ in } E^{J+1} : \pi'' \cdot \bar{\beta} = 1 \text{ and } |\pi'' - \pi'| < \rho\}$$

is a subset of A . If a distribution is not regular, then it (and its quasidensity and coordinate vector) will be called *irregular*. Equivalently, a distribution with quasidensity $g_{\pi'}$ is regular if and only if $Z^T(\pi' - \pi^0)$ is an interior point of B .

The uniform distribution and most distributions expected in applications are regular. However, if the pattern likelihood function $\ell(u^*, t)$ is unimodal with maximum at t_0 in $[c,d]$, then the unit step function at t_0 is an irregular distribution on $[c,d]$. The first result shows that some distributions are regular, i.e., that B is J dimensional.

III.1 If G is a distribution on $[c,d]$ with a positive quasidensity and if for $c < t < d$ the quasidensity of G evaluated at t equals the derivative of G at t then G is regular. In particular, the uniform distribution is regular.

Proof: If $\frac{d}{dt} G(t) = \sum_0^J \pi'_j h_j(t) = g_{\pi'}(t) > 0$ for $c < t < d$, and $g_{\pi'}$ is also positive at c and d , then $\min_t g_{\pi'}(t) > 0$. If $|\pi'' - \pi'|_{\infty} = \max_j |\pi''_j - \pi'_j|$ is sufficiently small, then $|g_{\pi'' - \pi'}(t)| \leq 1/2 \min_t g_{\pi'}(t)$. Consequently for sufficiently small $|\pi'' - \pi'|_{\infty}$

$$\begin{aligned} g_{\pi''}(t) &= g_{\pi'}(t) + g_{\pi'' - \pi'}(t) \\ &\geq \min_t g_{\pi'}(t) - \max_t |g_{\pi'' - \pi'}(t)| \end{aligned}$$

>0 .

Since $\|\cdot\|_2$ and $\|\cdot\|_\infty$ determine the same topology on E^{J+1} , for some $\rho_0 > 0$, $\|\pi'' - \pi'\|_2 < \rho_0$ implies $g_{\pi''}(t) > 0$ for t in $[c, d]$.

To show $\|\pi'' - \pi'\|_2 < \rho_0$ and $\pi'' \cdot \bar{\beta} = 1$ imply π'' is in A it suffices to show that these assumptions imply that the function \bar{G}

$$\bar{G}(x) = \begin{cases} 0 & \text{if } c < x \\ \int_c^x g_{\pi''}(t) dt & \text{if } c \leq x \leq d \\ 1 & \text{if } d < x \end{cases}$$

is a distribution on $[c, d]$ and $g_{\pi''}$ is its quasidensity. The only nontrivial step in verifying that \bar{G} is a distribution on $[c, d]$ is showing $\bar{G}(d) = 1$. Since for all t , $\sum_u \ell(u^*, t) = 1$, $\bar{G}(d) = \int_c^d g_{\pi''}(t) dt = \int_c^d \sum_u \ell(u^*, t) g_{\pi''}(t) dt = \sum_j \langle \sum_u \ell(u^*, \cdot), h_j \rangle \pi_j'' = \bar{\beta} \cdot \pi'' = 1$. Consequently, \bar{G} is a distribution on $[c, d]$, $g_{\pi''}$ is the restriction of its probability density to $[c, d]$, $g_{\pi''}$ is its quasidensity (by the uniqueness of quasidensities in I.1), and π'' is in A . In particular, since $1 = \sum_u \ell(u^*, t)$ is in the CS, the uniform density $g(t) = [d-c]^{-1}$ is in the CS, and the uniform distribution is regular. //

The next result gives examples of regular distributions that do not have continuous densities. It shows how to approximate any distribution on $[c, d]$ with a regular distribution.

III.2 If $t_0 < t_1 \dots < t_J$ are in $[c, d]$ and the vectors $\pi^i = (h_j(t_i))$ are linearly independent then for any vector α in E^{J+1} if

$$\alpha_j > 0 \quad j = 0, \dots, J$$

$$\sum_j \alpha_j = 1$$

then the discrete distribution function

$$G(t) = \sum_{(i: t_i \leq t)} \alpha_i$$

is regular. More generally if $G_0, G_1, \dots, G_i, \dots, G_J$ are distributions on $[c, d]$ with quasidensities g_i and the π^i are linearly independent, then $\alpha > 0$ and $\sum \alpha_i = 1$ imply that $\sum \alpha_i G_i$ is a regular distribution on $[c, d]$.

Proof: Clearly each for vector $\alpha > 0$ such that $\sum \alpha_i = 1$ the convex combination $\sum \alpha_i G_i(\cdot)$ of distributions G_i on $[c, d]$ is a distribution on $[c, d]$. Let $\alpha^0 > 0$ be the vector of coefficients of any such combination. A value of ρ will be computed to show $\sum \alpha_i^0 \pi^i$ is regular. For any α

$$\begin{aligned} \left| \sum_i \alpha_i^0 \pi^i - \sum_i \alpha_i \pi^i \right|^2 &= \sum_{ij} (\alpha_i^0 - \alpha_i) \pi^i \pi^j{}^T (\alpha_j^0 - \alpha_j) \\ &= (\alpha^0 - \alpha)^T Q (\alpha^0 - \alpha) \end{aligned}$$

for positive definite or positive semidefinite Q . Since the π^i are linearly independent, Q is definite. Consequently $\left| \sum_i \alpha_i^0 \pi^i - \sum_i \alpha_i \pi^i \right| \geq |\alpha^0 - \alpha| \epsilon^{1/2}$, where $\epsilon > 0$ is the smallest eigenvalue of Q . Since the $J+1$ linearly independent π^i form a basis for E^{J+1} , and for any π' in E^{J+1} , there is a unique α' in E^{J+1} such that $\pi' = \sum \alpha'_i \pi^i$. If

$$\left| \sum_i \alpha_i^0 \pi^i - \pi' \right| < \frac{1}{2} \epsilon^{1/2} \min_i (\alpha_i^0)$$

then

$$\begin{aligned} |\alpha^0 - \alpha'| &< \frac{1}{2} \min_i (\alpha_i^0), \\ \max_i |\alpha_i^0 - \alpha'_i| &< \frac{1}{2} \min_i (\alpha_i^0), \end{aligned}$$

and

$$0 < \alpha'_i \text{ for } i = 0, 1, \dots, J.$$

Furthermore, if $\pi' \cdot \bar{\beta} = 1$, then

$$\begin{aligned}
 1 &= (\sum_i \alpha'_i \pi^i) \cdot \bar{\beta} \\
 &= \sum_i \alpha'_i (\pi' \cdot \bar{\beta}) \\
 &= \sum_i \alpha'_i .
 \end{aligned}$$

Thus for $\rho = \frac{1}{2} \epsilon^{1/2} \min(\alpha_i^0)$,

$$|\sum_i \alpha_i^0 \pi^i - \pi'| < \rho \text{ and } \pi' \cdot \bar{\beta} = 1$$

imply $g_{\pi'}$ is the quasidensity of some distribution on $[c,d]$, i.e. π' is in A and $\sum_i \alpha_i^0 G_i$ is regular. //

The following result helps to visualize irregular distributions.

III.3 If π' is regular and π'' is in A then for sufficiently small positive ϵ , $\pi'' + (1+\epsilon)(\pi' - \pi'')$ is also in A .

Thus to obtain an irregular point one starts with any two points of A , π^0 and $\pi' \neq \pi^0$. Since A is convex, closed and bounded one can move through A along the ray from π^0 through π' to a point $\pi^0 + k(\pi' - \pi^0)$ in A such that $\pi^0 + k'(\pi' - \pi^0)$ is not in A for any $k' > k$. Thus $\pi^0 + k(\pi' - \pi^0)$ can not be regular.

Proof: $\bar{\beta} \cdot [\pi'' + (1+\epsilon)(\pi' - \pi'')] = 1 + (1+\epsilon)(1-1) = 1$ and, for sufficiently small positive ϵ , $|\pi' - (\pi'' + (1+\epsilon)(\pi' - \pi''))| = \epsilon |\pi' - \pi''| < \rho$. //

Irregular distributions can be obtained from the many unimodal functions (e.g. most likelihood functions) in the CS too.

III.4 If f is a function in the CS with a unique maximizer t_0 , i.e., if $f(t) \geq f(t_0)$ implies $t = t_0$ for all t in $[c,d]$, then the unit step function at t_0 is irregular.

Proof: Since f is a linear combination of likelihood functions for any distribution function on $[c,d]$, $\int_c^d f(t) dG(t) = \langle f, g \rangle$ where g is the

quasidensity of G . In particular for $g_{\pi(t_0)}$ = the quasidensity of the unit step at t_0 , $\langle f, g_{\pi(t_0)} \rangle = f(t_0)$. Note that if G is the distribution of a random variable ω then $\langle f, g \rangle = E[f(\omega)]$. Thus for every quasidensity π' in A , $\langle f, g_{\pi'} \rangle \leq \max_t f(t) = f(t_0)$. In particular for $g_{\pi(t_1)}$ the quasidensity of the unit step at $t_1 \neq t_0$,

$$\langle f, g_{\pi(t_1)} \rangle = f(t_1) < f(t_0) = \langle f, g_{\pi(t_0)} \rangle.$$

Consequently for any $\epsilon > 0$, for $\pi(\epsilon) = \pi(t_1) + (1+\epsilon)[\pi(t_0) - \pi(t_1)]$

$$\langle f, g_{\pi(\epsilon)} \rangle = \langle f, g_{\pi(t_0)} \rangle + \epsilon[\langle f, g_{\pi(t_0)} \rangle - \langle f, g_{\pi(t_1)} \rangle]$$

$$= f(t_0) + \epsilon[f(t_0) - f(t_1)] > f(t_0).$$

Thus $g_{\pi(\epsilon)}$ cannot be the quasidensity of any distribution on $[c, d]$ and from III.3, the unit step at t_0 is not regular. //

In fact a stronger result can be proven. It can be shown that the unit step at a maximizer corresponds to a point π' in A situated like a vertex of a polyhedron or a boundary point of an ellipsoid: If $\epsilon > 0$ and π'' is also in A , then $\pi' + (1+\epsilon)(\pi'' - \pi')$ is not in A . In other words, π' is not an interior point of the intersection of A and any line through π' .

III.5 If f is a function in the CS with unique maximizer t_0 , then

$\pi(t_0)$ is not an interior point of the intersection of A and any line through π' .

Proof: Let G be any distribution on $[c, d]$ other than the unit step at t_0 . Let $[c', d']$ be any closed subinterval of $[c, d]$ not containing t_0 such that $\int_{c'}^{d'} dG(t) > 0$. Then for $g_{\pi'}$, equal to the quasidensity of G ,

$$\langle f, g_{\pi'} \rangle = \int_{c'}^{d'} f(t) dG(t) + \int_{t \text{ not in } [c', d']} f(t) dG(t)$$

$$\begin{aligned}
&\leq \max_{c' \leq t \leq d'} f(t) \int_{c'}^{d'} dG(t) + f(t_0) \left[1 - \int_c^d dG(t) \right] \\
&< f(t_0) \int_{c'}^{d'} dG(t) + f(t_0) \left[1 - \int_c^d dG(t) \right] \\
&= f(t_0) = \langle f, g_{\pi(t_0)} \rangle .
\end{aligned}$$

From the argument in III.4 applied to $\pi(\epsilon) = \pi' + (1+\epsilon)[\pi(t_0) - \pi']$ it follows that for all $\pi' \neq \pi(t_0)$ in A for $\epsilon > 0$, $\pi(\epsilon)$ is not in A . //

Not every irregular distribution is mapped to one of these points that "stick out" from A like a vertex or a point of positive Gaussian curvature on a boundary. By taking linear combinations of unimodal functions in the CS one can construct bimodal and multimodal functions with modes of equal height. The reasoning used in III.4 and III.5 can then be used to show that A has edges and faces too.

III.6 Let f be a function in the CS having modes of equal height at

$t_0 < t_1 < \dots < t_m$ so that $f(t_0) = f(t_1) = \dots = f(t_m)$ and for t in $[c, d] \setminus \{t_0, t_1, \dots, t_m\}$, $f(t) < f(t_0)$. Let G_0, G_1, \dots, G_m denote the unit step functions at t_0, t_1, \dots, t_m . Then for any positive numbers α_i such that $\sum_{i=0}^m \alpha_i = 1$, the distribution $\sum \alpha_i G_i$ is irregular.

Proof: Let $g_{\pi(s)}$ be the quasidensity of the unit step function at s in $[c, d] \setminus \{t_0, t_1, \dots, t_m\}$. Then $\langle f, g_{\pi(s)} \rangle = f(s) < f(t_0) = \sum_{i=0}^m \alpha_i f(t_i)$. Thus for $g_{\pi(t_i)}$ equal to the quasidensity of the unit step at t_i and

$$\pi(\epsilon) = \pi(s) + (1+\epsilon)[\sum \alpha_i \pi(t_i) - \pi(s)]$$

$\epsilon > 0$ implies $\langle f, g_{\pi(\epsilon)} \rangle = f(t_0) + \epsilon[f(t_0) - f(s)] > f(t_0)$. Thus $\sum \alpha_i G_i$ is irregular. //

The following corollary reconciles the apparent contradiction between

III.2 and III.6 .

III.7 If for f in the CS there are more than J numbers t satisfying $f(t) \geq f(s)$ for all s in $[c,d]$ then the quasidensities of the unit step distributions at these numbers are linearly dependent.

Two ability distributions are called *equivalent* if the probability distribution of any function of the item scores does not depend on which of the two ability distributions is used to compute the distribution. Thus, if F and G are equivalent, then data cannot be used to determine which of the two distributions is correct. A necessary and sufficient condition for two distributions on $[c,d]$ to be equivalent is that they have the same quasidensity (Levine, 1989). We will show that for logistic models all distributions are equivalent to discrete distributions. The result is valid for models with item response functions P_i such that for each t there is a power series for P_i that converges absolutely in some neighborhood of t .

III.8 If the constant functions are the only functions in the CS that are constant on some nonempty open subset of $[c,d]$, then every distribution on $[c,d]$ is equivalent to a distribution with at most $J+1$ points of increase.

Proof: It is sufficient to prove that there are finitely many points of increase because A has been shown to be J dimensional. It is sufficient to limit attention to irregular distributions because if π' is not on the boundary ∂A of A then compactness of A implies that for any π^1 in ∂A we can choose $t > 1$ such that $\pi^2 = \pi^1 + t(\pi' - \pi^1)$ is also on ∂A . The equation

$$\pi' = \frac{t-1}{t} \pi^1 + \frac{1}{t} \pi^2$$

shows that π' corresponds to a probability mixture of distributions mapped to the boundary of A .

Let G be a distribution on $[c,d]$ with quasidensity $g_{\pi'}$ for π' on ∂A . Let $H = \{x \text{ in } E^{J+1} : n \cdot x = c\}$ be a hyperplane in E^{J+1} containing π' and no points of the interior of A . Without loss of generality it can be assumed that $x \text{ in } A$ implies $n \cdot x \leq c$ because $n \cdot x = c$ implies $(-n) \cdot x = -c$. It follows that among distributions on $[c,d]$ G is a maximizer of $\int_c^d \sum_j n_j h_j(t) dG(t) = n \cdot \pi'$. Either $g_n(\cdot) = \sum n_j h_j(\cdot)$ is constant on some subinterval of $[c,d]$ (and therefore constant) or there are only finitely many numbers t such that $g_n(t) \geq g_n(s)$ for all s in $[c,d]$. g_n cannot be constant for otherwise $n \cdot \pi'' = \int_c^d g_n(t) \cdot g_{\pi''}(t) dt = c$ would be independent of π'' and A would have no interior points. Thus for finitely many numbers $c \leq t_1 < \dots < t_K \leq d$ $g_n(t_k) = \max_{[c,d]} g_n(t)$. Thus a distribution \tilde{G} (such as G) maximizes $\int_c^d g_n(t) d\tilde{G}(t)$ if and only if it is equivalent to

$$\sum_{k=1}^K \alpha_k F_k(\cdot)$$

where F_k is the unit step at t_k for some positive numbers α_k such that $\sum \alpha_k = 1$. //

Section Four

Asymptotic Normality for Regular Distributions

In this section it will be shown that if the ability distribution is regular then $\hat{\pi}_N$, the maximum likelihood estimate of π , is asymptotically normal. A formula is derived for the asymptotic dispersion matrix.

Throughout this section the distribution for θ is assumed to be regular. Throughout this section let ρ be a fixed positive number chosen so that the intersection of the open ball with radius ρ centered at π

$$\{\pi' \text{ in } E^{J+1} : |\pi' - \pi| < \rho\}$$

and the hyperplane

$$\{\pi' \text{ in } E^{J+1} : \pi' \cdot \bar{\beta} = 1\}$$

is a subset of A . As in Section III.1, let z^1, z^2, \dots, z^J be any orthonormal basis for the annihilator of $\bar{\beta}$ and let $Z = [z^1, z^2, \dots, z^J]$ be the $(J+1) \times J$ matrix formed from the z 's so that $\pi' \rightarrow ZZ^T \pi'$ is the orthogonal projection onto the annihilator of $\bar{\beta}$.

To describe the asymptotic behavior of $\hat{\pi}_N$ we will need the information matrix, i.e. the matrix I of expected second derivatives with typical entry I_{ij}

$$\begin{aligned} I_{ij} &= -E \frac{\partial^2}{\partial \pi'_i \partial \pi'_j} \log P_{\pi'}(u) \Big|_{\pi'=\pi} \\ &= - \sum_u^* P_{\pi}(u^*) \frac{\partial^2}{\partial \pi'_i \partial \pi'_j} \log \pi' \cdot \beta(u^*) \Big|_{\pi'=\pi} \\ &= \sum_u^* P_{\pi}(u^*) \frac{\beta_i(u^*) \beta_j(u^*)}{[\pi \cdot \beta(u^*)]^2} \end{aligned}$$

Thus I can be written in the form $\sum_u^* w(u^*) \beta(u^*) \beta(u^*)^T$ and II.1 can be used to show it is non-singular. Since the columns of Z are independent, II.1 also implies that for any positive weights $w_{\nu} = w(u_{\nu}^*)$ the matrix

$\sum_{\nu=1}^{2^n} w_{\nu} Z^T \beta(u_{\nu}^*) \beta^T(u_{\nu}^*) Z$ is non-singular. In particular, $Z^T I Z$ is non-singular.

Since $\hat{\pi}_N$ almost surely converges to π , with probability one $\hat{\pi}_N$ will eventually be within ρ of π . For $\hat{s}_N = Z^T(\hat{\pi}_N - \pi)$ the mle can be written

$$\hat{\pi}_N = \pi + Z \hat{s}_N.$$

Since $|\hat{\pi}_N - \pi| = |\hat{s}_N|$, almost surely $|\hat{s}_N| < \rho$ for sufficiently large N . Defining M_N on open $\{s \text{ in } E^J : |s| < \rho\}$ by $M_N(s) = L_N[\pi + Zs]$ it follows that \hat{s}_N maximizes M_N and consequently must satisfy the equations

$$0 = \frac{\partial}{\partial s_j} M_N(s) \quad j = 1, 2, \dots, J.$$

In fact, \hat{s}_N almost surely eventually is the only solution of the equations with length less than ρ because with probability one, for sufficiently large N $J+1$ patterns with linearly independent β 's will be sampled, and this implies that the Hessian matrix evaluated at $|s| < \rho$

$$\partial^2 M_N(s) = -Z^T \left[\sum_{a=1}^N \beta(u_a^*) \beta^T(u_a^*) P_{\pi + Zs}(u_a^*)^{-2} \right] Z$$

is definite.

The asymptotic distribution of $\hat{\pi}_N$ is obtained with Taylor's formula applied to the gradient of M_N .

IV.2 (Asymptotic distribution of the maximum likelihood estimate). Let z^1, \dots, z^J be any orthonormal basis for $\text{Nul}(\bar{\beta})$ and $Z = [z^1, \dots, z^J]$. If the ability distribution is regular then $\hat{\pi}_N$ converges in distribution to

$$\pi + N^{-1/2} Z n$$

where n is multinormal with zero mean and covariance matrix $(Z^T I Z)^{-1}$.

Proof: Since $\hat{s}_N = Z^T(\hat{\pi}_N - \pi)$ implies that $Z\hat{s}_N = \hat{\pi}_N - \pi$, the theorem can be proven by showing that $N^{1/2}\hat{s}_N$ converges in distribution to multinormal n . This will be done by showing that for any non-zero J -vector t , the random variable $N^{1/2}t \cdot \hat{s}_N$ is asymptotically normal with mean zero and variance $t^T(Z^T I Z)^{-1}t$.

Each component of the gradient of M_N

$$\frac{\partial}{\partial s_k} M_N(s) = \sum_{a=1}^N [P_{\pi+Zs}(u_a^*)]^{-1} \beta(u_a^*) \cdot z^k \quad k = 1, \dots, J$$

is defined and has continuous partial derivatives of order two for $|s| < \rho$.

Thus when $|\hat{s}_N|$ is less than ρ there will be some $\epsilon_{N,k}$, $0 < \epsilon_{N,k} < 1$,

such that for each $k \leq J$

$$\begin{aligned} 0 &= \frac{\partial}{\partial s_k} M_N(0) + \sum_i \frac{\partial}{\partial s_i} \frac{\partial}{\partial s_k} M_N(0) \hat{s}_{N,i} + \frac{1}{2} \sum_i \left[\sum_j \frac{\partial^2}{\partial s_i \partial s_j} \frac{\partial}{\partial s_k} M_N(\epsilon_{N,k} \hat{s}_N) \hat{s}_{N,j} \right] \hat{s}_{N,i} \\ &= \frac{\partial}{\partial s_k} M_N(0) + \sum_i \left[\frac{\partial^2}{\partial s_i \partial s_k} M_N(0) + \frac{1}{2} \sum_j \frac{\partial^3}{\partial s_i \partial s_j \partial s_k} M_N(\epsilon_{N,k} \hat{s}_N) \hat{s}_{N,j} \right] \hat{s}_{N,i}. \end{aligned}$$

In matrix notation

$$0 = \partial M_N(0) + [\partial^2 M_N(0) + \frac{1}{2} C_N] \hat{s}_N$$

and

$$\left[-\frac{1}{N} \partial^2 M_N(0) - \frac{1}{N} \frac{1}{2} C_N \right] \hat{s}_N = \frac{1}{N} \partial M_N(0).$$

The right hand side of the last equation, being a mean of independent, identically distributed random vectors with zero expectation and covariance matrix $Z^T I Z$, is asymptotically normal with expectation 0 and covariance matrix $Z^T I Z/N$. Thus $N^{-1/2} \partial M_N(0)$ is asymptotically normal with mean zero and covariance matrix $Z^T I Z$. Since the summands in

$$-\frac{1}{N} \partial^2 M_N(0) = \frac{1}{N} \left(\sum_{a=1}^N [P_{\pi}(u_a^*)]^{-2} Z^T \beta(u_a^*) \beta(u_a^*)^T Z \right)$$

are independent and identically distributed, $-\frac{1}{N} \partial^2 M_N(0)$ converges almost surely to the non-singular matrix of expected values

$$\begin{aligned} E[P_{\pi}(u)^{-2} Z^T \beta(u) \beta(u)^T Z] &= \sum_{\nu=1}^{2^n} P_{\pi}(u_{\nu}^*) [P_{\pi}(u_{\nu}^*)]^{-2} Z^T \beta(u_{\nu}^*) \beta(u_{\nu}^*)^T Z \\ &= Z^T I Z . \end{aligned}$$

The k th row and i th column of C_N is

$$\begin{aligned} \sum_j \frac{\partial^3}{\partial s_i \partial s_j \partial s_k} M_N(\epsilon_N, k \hat{s}_N) \hat{s}_{N,j} \\ = \sum_{a=1}^N [P_{\pi+\epsilon_N, k}^{-3}(u_a^*)] \left[\beta(u_a^*) \cdot z^k \right] \left[\beta(u_a^*) \cdot z^i \right] \left[\beta(u_a^*) \cdot z \hat{s}_N \right] . \end{aligned}$$

Since \hat{s}_N converges almost surely to zero and the probabilities are bounded away from zero, the matrix $N^{-1} C_N$ converges almost surely to a matrix of zeros. Consequently the matrix D_N

$$D_N = \left(-\frac{1}{N} \partial^2 M_N(0) - \frac{1}{N} \frac{1}{2} C_N \right)$$

converges almost surely to non-singular $Z^T I Z$. Thus with probability one D_N is eventually non-singular and eventually both

$$\hat{s}_N = D_N^{-1} [N^{-1} \partial M_N(0)]$$

and

$$N^{1/2} t^T \hat{s}_N = t^T D_N^{-1} [N^{-1/2} \partial M_N(0)] .$$

Let $Y_N^T = \begin{cases} t^T D_N^{-1} & , \text{ if } D_N \text{ is non-singular} \\ 0^T & , \text{ otherwise} \end{cases}$

and $X_N = N^{-1/2} \partial M_N(0)$ so that $Y_N^T X_N$ almost surely eventually equals

$N^{1/2} t \cdot \hat{s}_N$. If it can be shown that $Y_N^T X_N$ converges in distribution to $t^T (Z^T I Z)^{-1} X$ where X is multivariate normal with covariance matrix $Z^T I Z$ then it follows that $Y_N^T X_N$ and $N^{1/2} t^T \hat{s}_N$ are asymptotically normal with variance $t^T (Z^T I Z)^{-1} (Z^T I Z) [t^T (Z^T I Z)^{-1}]^T = t^T (Z^T I Z)^{-1} t$, and the proof will be complete. $Y_N^T - t^T (Z^T I Z)^{-1} \xrightarrow{\text{wp1}} 0^T$ so $Y_N^T - t^T (Z^T I Z)^{-1} \xrightarrow{p} 0^T$. Since

X_N converges in distribution, $[Y_N^T - t^T(Z^T I Z)^{-1}]X_N \xrightarrow{d} 0$ and $\xrightarrow{p} 0$.

Finally, since $t^T(Z^T I Z)^{-1}X_N \xrightarrow{d} t^T(Z^T I Z)^{-1}X$,

$$Y_N^T X_N = [Y_N^T - t^T(Z^T I Z)^{-1}]X_N + t^T(Z^T I Z)^{-1}X_N \xrightarrow{d} t^T(Z^T I Z)^{-1}X. //$$

Acknowledgements

This research was supported by the Cognitive Science Program of the Office of Naval Research contract N00014-83K-0397, NR 150-518 and N00014-86K-0482, NR 4421546. I am indebted to Bruce Williams, Tim Davey, Charles Davis, Fritz Drasgow, Brian Junker, and Gary Thomasson for comments on earlier versions of this work. Conversations with J.O. Ramsay were also useful.

References

- Levine, M. V. (1985). The trait in latent trait theory. In D. J. Weiss (Ed.), Proceedings of the 1982 item response theory and computerized adaptive testing conference. Minneapolis, MN: University of Minnesota, Department of Psychology, Computerized Adaptive Testing Laboratory.
- Levine, M.V. (1989). Formula Scoring, Basic Theory and Applications. Report 89-1. Champaign, IL: Model-Based Measurement Laboratory, Department of Educational Psychology, University of Illinois.
- Levine, M.V. and Drasgow, F. (1988). Optimal appropriateness measurement. Psychometrika, 53, 161-176.
- Lord, F.M. (1970). Item characteristic curves estimated without knowledge of their mathematical form: A confrontation of Birnbaum's logistic model. Psychometrika, 35, 43-50.
- Samejima, F. (1981). Final report: Efficient methods of estimating the operating characteristics of item response categories and challenge to a new model for the multiple-choice item. Technical Report. Knoxville, Tennessee: Department of Psychology, University of Tennessee.
- Wald, A. (1949). Note on the consistency of the maximum likelihood estimate. Ann. Math. Statist., 20, 595-601.

University of Illinois/Levine

University of Illinois/Levine

Dr. Terry Ackerman
Educational Psychology
210 Education Bldg.
University of Illinois
Champaign, IL 61801

Dr. Robert Ahlers
Code N711
Human Factors Laboratory
Naval Training Systems Center
Orlando, FL 32813

Dr. James Algina
1403 Norman Hall
University of Florida
Gainesville, FL 32605

Dr. Erling B. Andersen
Department of Statistics
Studiestraede 6
1455 Copenhagen
DENMARK

Dr. Eva L. Baker
UCLA Center for the Study
of Evaluation
145 Moore Hall
University of California
Los Angeles, CA 90024

Dr. Isaac Bejar
Mail Stop: 10-R
Educational Testing Service
Rosedale Road
Princeton, NJ 08541

Dr. Menucha Dorenbaum
School of Education
Tel Aviv University
Ramat Aviv 69978
ISRAEL

Dr. Arthur S. Blawies
Code N712
Naval Training Systems Center
Orlando, FL 32813-7100

Dr. Bruce Bloxom
Defense Manpower Data Center
99 Pacific St.
Suite 155A
Monterey, CA 93943-3231

Dr. Stanley Collier
Office of Naval Technology
Code 222
800 N. Quincy Street
Arlington, VA 22217-5000

Dr. Hans F. Crombag
Faculty of Law
University of Limburg
P.O. Box 616
Maastricht
The NETHERLANDS 6200 MD

Dr. Timothy Davey
American College Testing Program
P.O. Box 168
Iowa City, IA 52243

Dr. C. M. Dayton
Department of Measurement
Statistics & Evaluation
College of Education
University of Maryland
College Park, MD 20742

Dr. Ralph J. DeAysla
Measurement, Statistics,
and Evaluation
Benjamin Bldg., Rm. 4112
University of Maryland
College Park, MD 20742

Dr. Dattprasad Divgi
Center for Naval Analysis
4401 Ford Avenue
P.O. Box 16268
Alexandria, VA 22302-0268

Dr. Hui-Ki Dong
Bell Communications Research
6 Corporate Place
PYA-1K226
Piscataway, NJ 08854

Dr. Fritz Drasgow
University of Illinois
Department of Psychology
603 E. Daniel St.
Champaign, IL 61820

Dr. R. Darrell Bock
Educational Psychology
NORC
6030 South Ellis
Chicago, IL 60637

Cdt. Arnold Bohrer
Sectie Psychologisch Onderzoek
Rekruterings-En Selectiecentrum
Kwartier Koningen Astrid
Bruijnstraat
1120 Brussels, BELGIUM

Dr. Robert Breau
Code 7B
Naval Training Systems Center
Orlando, FL 32813-7100

Dr. Robert Brennan
American College Testing
Programs
P. O. Box 168
Iowa City, IA 52243

Dr. John B. Carroll
409 Elliott Rd., North
Chapel Hill, NC 27514

Dr. Robert M. Carroll
Chief of Naval Operations
OP-01B2
Washington, DC 20350

Dr. Raymond E. Christal
UES LAMP Science Advisor
AFHRL/MOEL
Brooks AFB, TX 78235

Dr. Norman Cliff
Department of Psychology
Univ. of So. California
Los Angeles, CA 90089-1051

Director,
Manpower Support and
Readiness Program
Center for Naval Analysis
2000 North Beauregard Street
Alexandria, VA 22311

Defense Technical
Information Center
Cameron Station, Bldg 5
Alexandria, VA 22314
Attn: TC
(12 Copies)

Dr. Stephen Dunbar
224B Lindquist Center
for Measurement
University of Iowa
Iowa City, IA 52242

Dr. James A. Earles
Air Force Human Resources Lab
Brooks AFB, TX 78235

Dr. Kent Eaton
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Susan Ebbertson
University of Kansas
Psychology Department
426 Fraser
Lawrence, KS 66045

Dr. George Englehard, Jr.
Division of Educational Studies
Emory University
210 Fishburne Bldg.
Atlanta, GA 30322

Dr. Benjamin A. Fairbank
Performance Metrics, Inc.
5825 Callaghan
Suite 225
San Antonio, TX 78228

Dr. P-A. Federico
Code 51
NPRDC
San Diego, CA 92152-6800

Dr. Leonard Feldt
Lindquist Center
for Measurement
University of Iowa
Iowa City, IA 52242

Dr. Richard L. Ferguson
American College Testing
P.O. Box 168
Iowa City, IA 52243

Dr. Gerhard Fischer
Liebiggasse 5/3
A 1010 Vienna
AUSTRIA

Dr. Myron Fischl
U.S. Army Headquarters
DAPE-MRR
The Pentagon
Washington, DC 20310-0300

Prof. Donald Fitzgerald
University of New England
Department of Psychology
Armidale, New South Wales 2351
AUSTRALIA

Mr. Paul Foley
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Alfred R. Fragly
AFOSR/NL, Bldg. 410
Bolling AFB, DC 20332-6448

Dr. Robert D. Gibbons
Illinois State Psychiatric Inst.
Rm 529H
1601 W. Taylor Street
Chicago, IL 60612

Dr. Janice Gifford
University of Massachusetts
School of Education
Amherst, MA 01003

Dr. Robert Glaser
Learning Research
& Development Center
University of Pittsburgh
3939 O'Hara Street
Pittsburgh, PA 15260

Dr. Bert Green
Johns Hopkins University
Department of Psychology
Charles & 34th Street
Baltimore, MD 21218

Mr. Dick Hoshaw
OP-135
Arlington Annex
Room 2834
Washington, DC 20350

Dr. Lloyd Humphreys
University of Illinois
Department of Psychology
603 East Daniel Street
Champaign, IL 61820

Dr. Steven Hunka
3-104 Educ. N.
University of Alberta
Edmonton, Alberta
CANADA T6G 2G5

Dr. Huynh Huynh
College of Education
Univ. of South Carolina
Columbia, SC 29208

Dr. Robert Jannerone
Elec. and Computer Eng. Dept.
University of South Carolina
Columbia, SC 29208

Dr. Douglas H. Jones
Thatcher Jones Associates
P.O. Box 6540
10 Trafalgar Court
Lawrenceville, NJ 08648

Dr. Brian Junker
University of Illinois
Department of Statistics
101 Illinois Hall
725 South Wright St.
Champaign, IL 61820

Dr. Milton S. Katz
European Science Coordination
Office
U.S. Army Research Institute
Box 65
FPO New York 09510-1500

Prof. John A. Keats
Department of Psychology
University of Newcastle
N.S.W. 2308
AUSTRALIA

DORNIER GMBH
P.O. Box 1420
D-7990 Friedrichshafen 1
WEST GERMANY

Prof. Edward Haertel
School of Education
Stanford University
Stanford, CA 94305

Dr. Ronald K. Hambleton
University of Massachusetts
Laboratory of Psychometric
and Evaluative Research
Hills South, Room 152
Amherst, MA 01003

Dr. Delwyn Harnisch
University of Illinois
51 Gerty Drive
Champaign, IL 61820

Dr. Grant Henning
Senior Research Scientist
Division of Measurement
Research and Services
Educational Testing Service
Princeton, NJ 08541

Ms. Rebecca Hetter
Navy Personnel R&D Center
Code 63
San Diego, CA 92152-6800

Dr. Paul W. Holland
Educational Testing Service, 21-1
Rosedale Road
Princeton, NJ 08541

Prof. Lutz F. Horneke
Institut für Psychologie
RWTH Aachen
Jaegerstrasse 17/19
D-5100 Aachen
WEST GERMANY

Dr. Paul Horst
677 G Street, #184
Chula Vista, CA 92010

Dr. G. Gage Kingsbury
Portland Public Schools
Research and Evaluation Department
501 North Dixon Street
P. O. Box 3107
Portland, OR 97209-3107

Dr. William Koch
Box 7246, Meas. and Eval. Ctr.
University of Texas-Austin
Austin, TX 78703

Dr. Leonard Kreeker
Navy Personnel R&D Center
Code 62
San Diego, CA 92152-6800

Dr. Jerry Lehnus
Defense Manpower Data Center
Suite 400
1600 Wilson Blvd
Reston, VA 22209

Dr. Thomas Leonard
University of Wisconsin
Department of Statistics
1210 West Dayton Street
Madison, WI 53705

Dr. Michael Levine
Educational Psychology
210 Education Bldg.
University of Illinois
Champaign, IL 61801

Dr. Charles Lewis
Educational Testing Service
Princeton, NJ 08541-0001

Dr. Robert L. Linn
Campus Box 249
University of Colorado
Boulder, CO 80309-0249

Dr. Robert Lockman
Center for Naval Analysis
4401 Ford Avenue
P.O. Box 16268
Alexandria, VA 22302-0268

Dr. Frederic M. Lord
Educational Testing Service
Princeton, NJ 08541

University of Illinois/Levine

University of Illinois/Levine

Dr. George B. Macready
Department of Measurement
Statistics & Evaluation
College of Education
University of Maryland
College Park, MD 20742

Dr. Gary Marco
Stop 31-E
Educational Testing Service
Princeton, NJ 08451

Dr. James R. McBride
The Psychological Corporation
1250 Sixth Avenue
San Diego, CA 92101

Dr. Clarence C. McCormick
HQ, USMEPCOM/MEPC
2500 Green Bay Road
North Chicago, IL 60064

Mr. Christopher McCusker
University of Illinois
Department of Psychology
603 E. Daniel St.
Champaign, IL 61820

Dr. Robert McKinley
Law School Admission Services
Box 40
Newtown, PA 18940

Dr. James McMichael
Technical Director
Navy Personnel R&D Center
San Diego, CA 92152-6800

Mr. Alan Mead
c/o Dr. Michael Levine
Educational Psychology
210 Education Bldg.
University of Illinois
Champaign, IL 61801

Dr. Robert Mislavsky
Educational Testing Service
Princeton, NJ 08541

Dr. William Montague
NPRDC Code 13
San Diego, CA 92152-6800

Dr. James B. Olsen
MICA Systems
1875 South State Street
Orem, UT 84058

Office of Naval Research,
Code 1142CS
800 N. Quincy Street
Arlington, VA 22217-5000
(6 Copies)

Office of Naval Research,
Code 125
800 N. Quincy Street
Arlington, VA 22217-5000

Assistant for MPI Research,
Development and Studies
OP 0187
Washington, DC 20370

Dr. Judith Orasanu
Basic Research Office
Army Research Institute
5001 Eisenhower Avenue
Alexandria, VA 22333

Dr. Jesse Orlansky
Institute for Defense Analyses
1801 N. Beauregard St.
Alexandria, VA 22311

Dr. Peter J. Pashley
Educational Testing Service
Rosedale Road
Princeton, NJ 08541

Wayne H. Patience
American Council on Education
GED Testing Service, Suite 20
One Dupont Circle, NW
Washington, DC 20036

Dr. James Paulson
Department of Psychology
Portland State University
P.O. Box 751
Portland, OR 97207

Dept. of Administrative Sciences
Code 54
Naval Postgraduate School
Monterey, CA 93943-5026

Ms. Kathleen Moreno
Navy Personnel R&D Center
Code 62
San Diego, CA 92152-6800

Headquarters Marine Corps
Code MPJ-20
Washington, DC 20380

Dr. Ratna Nandakumar
Dept. of Educational Studies
Willard Hall, Room 213
University of Delaware
Newark, DE 19716

Dr. W. Alan Nicewander
University of Oklahoma
Department of Psychology
Norman, OK 73071

Deputy Technical Director
NPRDC Code 01A
San Diego, CA 92152-6800

Director, Training Laboratory,
NPRDC (Code 05)
San Diego, CA 92152-6800

Director, Manpower and Personnel
Laboratory,
NPRDC (Code 06)
San Diego, CA 92152-6800

Director, Human Factors
& Organizational Systems Lab,
NPRDC (Code 07)
San Diego, CA 92152-6800

Library, NPRDC
Code P201L
San Diego, CA 92152-6800

Commanding Officer,
Naval Research Laboratory
Code 2627
Washington, DC 20390

Dr. Harold F. O'Neil, Jr.
School of Education - MPH 801
Department of Educational
Psychology & Technology
University of Southern California
Los Angeles, CA 90089-0031

Department of Operations Research,
Naval Postgraduate School
Monterey, CA 93940

Dr. Mark D. Reckase
ACT
P. O. Box 168
Iowa City, IA 52243

Dr. Malcolm Ree
AFHRL/MOHA
Brooks AFB, TX 78235

Mr. Steve Reiss
N660 Elliott Hall
University of Minnesota
75 E. River Road
Minneapolis, MN 55455-0344

Dr. Carl Ross
CHET-POCD
Building 90
Great Lakes NTC, IL 60088

Dr. J. Ryan
Department of Education
University of South Carolina
Columbia, SC 29208

Dr. Fumiko Samejima
Department of Psychology
University of Tennessee
310B Austin Peay Bldg.
Knoxville, TN 37916-0000

Mr. Drew Sands
NPRDC Code 62
San Diego, CA 92152-6800

Lowell Scheer
Psychological & Quantitative
Foundations
College of Education
University of Iowa
Iowa City, IA 52242

Dr. Mary Schratz
905 Orchid Way
Carlsbad, CA 92009

Dr. Dan Segall
Navy Personnel R&D Center
San Diego, CA 92152

Dr. W. Steve Sellman
OASD(HR&SL)
20269 The Pentagon
Washington, DC 20301

Dr. Kazuo Shigenasu
7-9-24 Kugenuma-Kaigan
Fujisawa 251
JAPAN

Dr. William Sims
Center for Naval Analysis
4401 Ford Avenue
P.O. Box 16268
Alexandria, VA 22302-0268

Dr. H. Wallace Sinako
Manpower Research
and Advisory Services
Smithsonian Institution
801 North Pitt Street, Suite 120
Alexandria, VA 22314-1713

Dr. Richard E. Snow
School of Education
Stanford University
Stanford, CA 94305

Dr. Richard C. Sorensen
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Judy Spray
ACT
P.O. Box 168
Iowa City, IA 52243

Dr. Martha Stocking
Educational Testing Service
Princeton, NJ 08541

Dr. Peter Stoleff
Center for Naval Analysis
4401 Ford Avenue
P.O. Box 16268
Alexandria, VA 22302-0268

Dr. William Stout
University of Illinois
Department of Statistics
101 Illini Hall
725 South Wright St.
Champaign, IL 61820

Dr. Ledyard Tucker
University of Illinois
Department of Psychology
603 E. Daniel Street
Champaign, IL 61820

Dr. David Vale
Assessment Systems Corp.
2233 University Avenue
Suite 440
St. Paul, MN 55114

Dr. Frank L. Vicino
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Howard Warner
Educational Testing Service
Princeton, NJ 08541

Dr. Ming-Mei Wang
Lindquist Center
for Measurement
University of Iowa
Iowa City, IA 52242

Dr. Thomas A. Warr
FAA Academy AAC934D
P.O. Box 25082
Oklahoma City, OK 73125

Dr. Brian Waters
HunARO
12505 Argyle Circle
Alexandria, VA 22314

Dr. David J. Weiss
N660 Elliott Hall
University of Minnesota
75 E. River Road
Minneapolis, MN 55455-0344

Dr. Ronald A. Weitzman
Box 146
Carnel, CA 93921

Major John Welsh
AFHRL/MOAN
Brooks AFB, TX 78223

Dr. Hariharan Swaminathan
Laboratory of Psychometric and
Evaluation Research
School of Education
University of Massachusetts
Amherst, MA 01003

Mr. Brad Symphon
Navy Personnel R&D Center
Code-131
San Diego, CA 92152-6800

Dr. John Tangney
AFOSR/ML, Bldg. 410
Bolling AFB, DC 20332-6448

Dr. Kikumi Tatsuka
CERL
252 Engineering Research
Laboratory
103 S. Mathews Avenue
Urbana, IL 61801

Dr. Maurice Tatsuka
220 Education Bldg
1310 S. Sixth St.
Champaign, IL 61820

Dr. David Thissen
Department of Psychology
University of Kansas
Lawrence, KS 66044

Mr. Thomas J. Thomas
Johns Hopkins University
Department of Psychology
Charles & 34th Street
Baltimore, MD 21218

Mr. Gary Thomaston
University of Illinois
Educational Psychology
Champaign, IL 61820

Dr. Robert Tsutakawa
University of Missouri
Department of Statistics
222 Math. Sciences Bldg.
Columbia, MO 65211

Dr. Douglas Wetzel
Code 51
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. Rand R. Wilcox
University of Southern
California
Department of Psychology
Los Angeles, CA 90089-1061

German Military Representative
ATTN: Wolfgang Wildgrube
Streitkräfteamt
D-5300 Bonn 2
4000 Brandynne Street, NW
Washington, DC 20016

Dr. Bruce Williams
Department of Educational
Psychology
University of Illinois
Urbana, IL 61801

Dr. Hilda Wing
MRC MH-176
2101 Constitution Ave.
Washington, DC 20418

Mr. John H. Wolfe
Navy Personnel R&D Center
San Diego, CA 92152-6800

Dr. George Wong
Biostatistics Laboratory
Memorial Sloan-Kettering
Cancer Center
1275 York Avenue
New York, NY 10021

Dr. Wallace Wulfeck, III
Navy Personnel R&D Center
Code 51
San Diego, CA 92152-6800

Dr. Kentaro Yamamoto
03-1
Educational Testing Service
Rosedale Road
Princeton, NJ 08541

1989/08/04

University of Illinois/Levine

Dr. Wendy Yen
CTB/McGraw Hill
Del Monte Research Park
Monterey, CA 93940

Dr. Joseph L. Young
National Science Foundation
Room 320
1800 G Street, N.W.
Washington, DC 20550

Mr. Anthony R. Zara
National Council of State
Boards of Nursing, Inc.
625 North Michigan Avenue
Suite 1544
Chicago, IL 60611